

Feature Article

An overview of heavy metal contamination in coastal sediments of Sri Lanka

A.M.N.M. Adikaram¹ and H.M.T.G.A. Pitawala²

¹Department of Physical Sciences, Faculty of Applied Sciences, South Eastern University, Sri Lanka

²Department of Geology, Faculty of Science, University of Peradeniya, Sri Lanka

Summary

Coastal sediments are often subjected to heavy metal contamination as they reside at the marginal environments of anthropological water releases. These sediments provide the habitat for marine aquatic life of seafood. Therefore, heavy metal contamination in coastal sediments is one of the major environmental concerns. The coastal belt of Sri Lanka is highly urbanized with high population (59% of Sri Lankan population) and is mostly depend on the sea. Therefore, anthropogenic inputs of pollutants in to the marine environments have been increased during last decades. Heavy metal contamination of the coastal sediments of Sri Lanka has been discussed by several studies, focusing on selected coastal regions. In this article, previous studies have been reviewed in order to understand the contamination status and ecological risks due to heavy metal accumulations. Despite the sampling location, the elemental distribution of coastal lagoonal sediments shows similar trend indicating the dominance of natural elemental sources rather the anthropogenic influences. The concentrations of As and Cr in sediments are high compared to the upper continental crust values which is a characteristic feature in possible source rocks, soils and sediments of Sri Lanka. The results of the previous studies indicate that coastal sediments are low to moderately contaminated but not at ecological risk. However, anthropogenic activities are highly variable across the coastal regions. Since limited results of the previous studies is not enough to get an overview of the heavy metal concentrations around the island, urgent need for a spatial and temporal geochemical database for coastal sediments of Sri Lanka is emphasized.

*Correspondence: maduryaa@gmail.com

Tel: +94718318140

ISSN 2235-9370 Print/ISSN 2235-9362 Online © 2017 University of Sri Jayewardenepura

Introduction

Environmental contaminants are in three categories: (a) stable trace elements (b) organic substances and (c) radionuclides (Tornero and d'Alcalà, 2014). United States Environmental Protection Agency (USEPA) has chosen the following trace elements including heavy metals such as Ag, As, Ba, Be, Cd, Co, Cr, Cu, Hg, Mn, Mo, Ni, Pb, Sb, Se and Zn that is toxic in desired excessive concentrations. The list has expanded later by several authors (McBride, 1995; Chen and Ma, 1998).

When a substance is above the level of background it is said to be as 'contaminated'. When it is adversely affected on biological communities, it is said to be as 'polluted' (Chapman, 2007). Heavy metal contamination in natural environments is a severe environmental problem due to their natural toxicity, persistence, vast sources and non-degradability (Wang et al., 2013). Heavy metals get into natural environments from effluents of industries, agriculture and aquaculture practices, and municipal and domestic wastes (Tornero and d'Alcalà, 2014). Marine and coastal environments have been paid more attention because they are the ultimate settings of heavy metal accumulation and human concern for seafood (Wang et al., 2005). Bioaccumulation, bio-magnification by food chains, destruction to organisms and ecological systems and consequently damage to ecological goods and services are the adverse effects of heavy metal contaminations (Newton et al., 2013; Tornero and d'Alcalà, 2014). Therefore, it is essential to monitor the present contamination status in marine and coastal environments.

Sri Lankan coastline is about 1920 km long and encompasses about 50 % of the Bay of Bengal (Katupotha, 1989). Different coastal features such as lagoons, estuaries, sediment bars and spits, dunes, wave cuts, arches and cliffs which are characterized by erosional and depositional processes are present around the island (Katupotha, 2007). The origin and evolution of coastal landforms of the country was activated during the Quaternary period and the sea level reaches its present status about 3000 yr BP (Ranasinghe et al., 2013). Several studies on different coastal landforms of Sri Lanka have been reviewed in this article to summarize the current status of coastal sediment contamination. This article reviews the existing literature on the concentrations of selected trace elements in the coastal regions of Sri Lanka and risk of contamination of the coastal ecosystems by them.

Coastal environment of Sri Lanka

The climate of Sri Lanka is tropical with diversified rainfalls from both southwest (SW) and northeast (NE) monsoons. In general, high rainfall is recorded during May to September due to SW monsoons and low rainfall during November to February due to NE monsoons. Altogether 103 main rivers are flowing to the coastal low lands from the central part of the country. The river discharges are timely changed with monsoonal rainfall effect. Huge sediment loads are carried through the river channels, and hence river channels are the basic sediment supplier for the coastal zones.

Large scale storm or cyclone has not been recorded during past decades except the tsunami occurred in 2004. The temperature ranges from 16 °C to 32 °C across the country. The lowest temperature recorded in the central highlands and highest temperatures are recorded in constantly southeastern and eastern coastal areas. The tidal effect of the island is micro-scale and hence no considerable effect on sedimentation in coastal zones except the regionally developing sea currents.

Many of the coastal zones of Sri Lanka are densely populated due to clear and shallow beaches, corals and hot sun. About 59% of the total population of Sri Lanka is associated to the coastal regions (Nayanananda, 2007). Most of the industrialized capital cities are located around the coastal margins of the country. Sea around the country receives nutrient rich currents, therefore other than point sources of contaminants non-point source such as storm water and dust can contribute for contamination. Fishing industry is famous in Sri Lanka and fish processing units are established around the coastal areas. Western and southern coastal regions are highly industrialized and densely populated. Thus, effluents that comes from industries such as petroleum refining, ship repairing, tires and thermal industries other than the huge municipal and domestic leachates are mixed with natural waters are ended up at the sea. Additionally, minor water based industries such as foundry, laundry, tannery, paper industries, service stations and 'batik' fabric die work are commonly observed in coastal low lands of Sri Lanka where tourists are attracted.

Levels of contaminants

In present article, heavy metals and As concentrations in sediments of seven main semi-enclosed marginal marine water bodies and capital city of Sri Lanka are summarized from published data. These recent publications are done by collecting surface sediment fractions of each semi-enclosed body. Sampling locations are Trincomalee bay, Batticaloa lagoon, Hambanthota lagoon, Galle harbor, Hikkaduwa lagoon, Beruwala harbor, Muthurajawela peat deposit and surface sediments of Colombo city (Dissanayake, 1987; Jayawardana et al., 2012; Young et al., 2014; Silva, 2015; Heath et al., 2016; Adikaram et al., 2017).

Geographical distribution of the concentrations of the selected heavy metals of Sri Lanka is shown in Fig. 1 (Dissanayake, 1987; Jayawardana et al., 2012; Young et al., 2014; Silva, 2015; Heath et al., 2016; Adikaram et al., 2017). Concentrations of Zn and Cr are significant in all natural lagoonal systems including Muthurajawela peat deposit with compared to the other elements. The two harbors show high Pb concentrations. Substantial enrichments of Zn and Cu are observed in the city sediments of the Colombo region.




Figure 1: Geographical distribution of heavy metal concentrations in coastal sediments of Sri Lanka. (After Dissanayake, 1987; Jayawardana et al., 2012; Young et al., 2014; Silva, 2015; Herath et al., 2016; Adikaram et al., 2017)

Average concentrations of As, Pb, Zn, Cu, Ni and Cr of the semi-enclosed water bodies of Sri Lanka, other published data in different regions of the world and international standards are summarized in Table 1. It is noted that the sediments of Sri Lankan coastal zones are not contaminated with compared to other compared marine environments of the world.

Table 1: Concentrations of heavy metals and As in coastal sediments of Sri Lanka and comparison with other coasts

Country	Location	Concentration (ppm)						Reference
		As	Pb	Zn	Cu	Ni	Cr	
Sri Lanka	Batticaloa lagoon	5 (3-10)*	21 (14-26)	87 (21-155)	34 (6-216)	18 (7-41)	93 (52-146)	Adikaram et al., 2017
	Trincomalee, inner harbour	7 (3-19)	23 (3-103)	49 (21-152)	13 (2-41)	16 (3-57)	73 (26-192)	Young et al., 2014
	Trincomalee, Thambalagam bay	10 (2-43)	14 (8-20)	31 (9-79)	6 (0-23)	11 (4-46)	50 (11-181)	Young et al., 2014
	Trincomalee, Koddiyar bay	4 (2-9)	15 (9-20)	51 (6-181)	6 (2-14)	13 (0-27)	96 (7-356)	Young et al., 2014
	Hikkaduwa lagoon	7 (3-11)	13 (8-17)	35 (8-55)	6 (2-9)	8 (0-20)	60 (45-84)	Jayawardana et al., 2012
	Hambanthota lagoon	3 (3-4)	14 (14-15)	47 (38-72)	9 (5-12)	11 (10-17)	72 (61-78)	Jayawardana et al., 2012
	Galle fishery harbour	nd	74 (61-87)	16 (1-68)	10 (7-16)	nd	nd	Silva (2015)
	Beruwala fishery harbour	nd	80 (37-102)	12 (0-50)	11 (5-15)	nd	nd	Silva (2015)
	Muthurajawela peat	nd	nd	67 (26-262)	20 (12-37)	nd	41 (20-84)	Dissanayake (1987)
Colombo city dust	nd	26 (0-76)	270 (53-625)	91 (28-319)	nd	80 (47-152)	Herath et al., (2016)	
India	Gulf of Mannar	nd	16	73	57	24	177	Jonathan et al., 2009
	Kalpakkam	nd	22	119	nd	53	118	Selvaraj et al., 2004
	Pondicherry	nd	33	52	48	20	334	Solali et al, 2013
	Gange's estuary	nd	29	71	26	32	67	Subramanian et al., 1988
	Krishna Estuary	nd	4	1482	59	149	174	Ramesh and Subramanium, 1988
	West Bengal	nd	17	74	36	34	37	Sarkar et al., 2004
Bangladesh	Chittagong coast,	nd	18	355	189	33	658	Hasan et al., 2013
Tunisia	Mediterranean sea	(13-36)	(65-152)	(47-546)	(8-29)	(9-31)	nd	Zohra and Habib, 2016
China	Bhohai bay	nd	(21-66)	(55-457)	(20-63)	(23-53)	(60-224)	Gao and Chen, 2012
SQG	TEC	10	36	121	32	23	43	McDonald et al., 1996
	PEC	33	128	459	149	49	111	McDonald et al., 1997

SQG: sediment quality guidelines, TEC: threshold effect level, PEC: probable effect level

The concentration of As and Cr in coastal sediments of Sri Lanka is high with compared to the upper continental crust (UCC) values (Fig. 1; Taylor and McLennan, 1985). The possible source country rock types are characterized in high concentrations of As and Cr and this might be the reason for the enrichments of such elements (Young et al., 2014).




Figure 2: UCC normalized elemental concentrations in coastal sediments of Sri Lanka (After Dissanayake, 1987; Jayawardana et al., 2012; Young et al., 2014; Silva, 2015; Heath et al., 2016; Adikaram et al., 2017).

Environmental and ecological risk assessment of coastal sediments

Average concentrations of six trace elements of coastal sediments of Sri Lanka are compared with reference values of Threshold Effect Concentration (TEC) and Probable Effect Concentrations (PEC) of consensus-based freshwater ecosystem Sediment Quality Guidelines (Table 1; MacDonald et al., 1996). Values in between TEC and PEC are rarely or occasionally make adverse effects on organisms whereas values above PEC make adverse effects on a wide range of organisms (MacDonald et al., 1996). The average concentrations of As of Sri Lankan coasts are below the threshold values of sediment quality guidelines. In addition, Beruwala and Galle harbors shows above Pb values for TEC. The concentrations of Zn, Cu and Cr of Colombo

city sediments show higher values for TEC. Except Muthurajawela peaty sediments, all sediment types have exceeded the TEC values for Cr. However, average concentrations of all sediment types of Sri Lankan coasts are below the PEC values suggesting the level of contamination is not much considerable.

Contamination factor (CF) is an elemental ratio of concentration of each metal in the sediment with concentration of those particular metals in the background (Hakanson, 1980). The contamination levels may be classified based on their intensities on a scale ranging from 1 to 6 (0 = none, 1 = none to medium, 2 = moderate, 3 = moderately to strong, 4 = strongly polluted, 5 = strong to very strong, 6 = very strong (Muller, 1969).

$$CF = \frac{CM_{Sediment}}{CM_{background}} \dots\dots\dots \text{(Equation1)}$$

Where, *CM* is the metal concentration.

The pollution load index (PLI) is calculated using the CF values that provide comparative means to measure the level of pollution in different locations (Tomlinson et al., 1980).

$$PLI = \sqrt[n]{CF_1 \times CF_2 \times CF_3 \times \dots \times CF_n} \dots\dots\dots \text{(Equation2)}$$

Where, CF is the contamination factor and n is the number of elements. If sediments are polluted $PLI > 1$ and for non-polluted $PLI < 1$ (Tomlinson et al., 1980).

The CF values and PLI values for average concentrations of semi-enclosed coastal bodies of Sri Lanka are given in Table 2. Contamination factor also indicates enrichments of As and Cr in Sri Lanka coasts. According to the PLI values, Colombo city sediments are significantly polluted. Additionally, Batticaloa lagoon and Trincomalee bay shows slightly polluted sediments whereas other locations are unpolluted with the global references. This is basically due to the high concentrations of Cr and As which is controlled by the country rocks.

Table 2: Contamination factor, pollution load index and potential ecological risk index of coastal sediments, Sri Lanka

Location	Contamination factor						PLI	PERI
	As	Pb	Zn	Cu	Ni	Cr		
Batticaloa lagoon	3.3	1.1	1.2	1.4	0.9	2.7	1.6	56
Trincomalee bay	3.3	1.1	1.2	1.4	0.9	2.7	1.6	56
Hikkaduwa lagoon	4.7	0.7	0.5	0.2	0.4	1.7	0.8	57
Hambanthota lagoon	2.0	0.7	0.7	0.4	0.6	2.1	0.9	33
Galle harbour	nd	3.7	0.2	0.4	nd	nd	0.7	21
Beruwala harbour	nd	4.0	0.2	0.4	nd	nd	0.7	22
Muthurajawela peat	nd	nd	0.9	0.8	nd	1.2	1.0	7
Colombo city	nd	1.3	3.8	3.6	nd	2.3	2.5	33

PLI: pollution load index

PERI: potential ecological risk index

High metal concentrations in sediments create adverse biological effects which can be measured by several defined approaches. To evaluate the adverse biological effect a simple method introduced by Hakanson (1980) was used. The potential ecological risk index (PERI) is specially proposed to assess the contamination levels of coastal sediments with respect to toxicity of some selected contaminants (Hakanson, 1980).

$$PERI = \sum E_r \dots\dots\dots (Equation 3)$$

$$E_r = T_r \times CF \dots\dots\dots (Equation 4)$$

Where, E_r is the ecological risk factor, T_r is the toxic response factor and CF is the contamination factor. The standard values for T_r are As, Pb, Zn, Cu, Ni and Cr are 10, 5, 1, 5, 5 and 2 respectively. The $PERI < 90$ is low potential ecological risk; $90 < PERI < 190$ is moderate ecological risk; $190 < PERI < 380$ is considerable ecological risk and $PERI > 380$ is a very high ecological risk. According to the calculated PERI values with published data, it can be suggested that the Sri Lankan coastal sediments have low potential ecological risk (Table 2).

Summery

In summary, based on the available data of Sri Lankan coastal sediments, it can be concluded that the sediments are slightly polluted and that level is not adversely affected to the ecological systems. The selection of geochemical background values in contamination status evaluations might be a reason for the pollution status of the present article which is basically depends on the possible source rocks. Regional trace metal concentrations are affected by the rock types, geomorphological conditions and geographical locations which are altered by physical and chemical weathering processes. Therefore, to determine the anthropogenic influences more accurately, a baseline for the sediment trace metal concentrations should be determined regionally. This will help to identify the anthropogenic causes of pollution and take necessary actions to control pollution to prevent the present coastal sediments.

References

- Adikaram, M., Pitawala, A., Ishiga, H., Jayawardana, D., (2017). Spatial distribution, enrichment, and source of environmentally important elements in Batticaloa lagoon, Sri Lanka. *Environmental Science and Pollution Research* 24(2), 2089-2099.
- Chapman, P. M. (2007). Determining when contamination is pollution—weight of evidence determinations for sediments and effluents. *Environment International* 33(4), 492-501.
- Chen, M., Ma, L. Q., (1998). Comparison of four USEPA digestion methods for trace metal analysis using certified and Florida soils. *Journal of Environmental Quality* 27(6), 1294-1300.
- Gao, X., Chen, C. T. A. (2012). Heavy metal pollution status in surface sediments of the coastal Bohai Bay. *Water research* 46(6), 1901-1911.
- Hakanson, L., (1980). An ecological risk index for aquatic pollution control. A sedimentological approach. *Water Research* 14, 975–1001.
- Hasan, A.B., Kabir, S., Selim Reza, A.H.M., Zaman, M.N., Ahsan, A., Rashid, M., (2013). Enrichment factor and geo-accumulation index of trace metals in sediments of the ship breaking area of Sitakund Upazilla (Bhatiary-Kumira), Chittagong, Bangladesh. *Journal of geochemical exploration* 125, 130–137.

- Herath, D., Pitawala, A., Gunatilake, J. (2016). Heavy metals in road deposited sediments and road dusts of Colombo Capital, Sri Lanka. *Journal of the National Science Foundation of Sri Lanka* 44(2).
- Jayawardana, D.T., Ishiga, H., Pitawala, H.M.T.G.A., (2012). Geochemistry of surface sediments in tsunami-affected Sri Lankan lagoons regarding environmental implications. *International journal of environmental science and technology* 9, 41–55.
- Jonathan, M.P., Ram Mohan, V., (2003). Heavy metals in sediments of the inner shelf off the Gulf of Mannar, Southeast coast of India. *Marine pollution bulletin* 46, 263–268.
- Katupotha, J., (1989). Coastal landforms during the Holocene epoch in Sri Lanka: are they comparable to those in Brazil and Venezuela? *International symposium on global changes in South America during the Quaternary, Sao Paulo, Brazil*, 188-192.
- Katupotha, K.N.J., (2007). (Coastal Landforms) *The National Atlas of Sri Lanka, Second Edition*, Survey Department of Sri Lanka. 40-42.
- MacDonald, D.D., Carr, R.S., Calder, F.D., Long, E.R., Ingersoll, C.G., (1996). Development and evaluation of sediment quality guidelines for Florida coastal waters. *Ecotoxicology* 5(4), 253–278.
- McBride, M. B., (1995). Toxic metal accumulation from agricultural use of sludge: are USEPA regulations protective?. *Journal of Environmental Quality* 24(1), 5-18.
- Muller, G. (1969). Index of Geoaccumulation in Sediments of the Rhine River. *GeoJournal* 2, 108-118.
- Nayanananda, O. K. (2007). *The Study of Economic Significance of Coastal Region of Sri Lanka in the Context of Environmental Changes of Pre and Post Tsunami*. Coast Conservation Department and The Ministry of Environment and Natural Resources.
- Newton, A., Icelly, J., Cristina, S., Brito, A., Cardoso, A. C., Colijn, F., ... Ivanova, K., (2014). An overview of ecological status, vulnerability and future perspectives of European large shallow, semi-enclosed coastal systems, lagoons and transitional waters. *Estuarine, Coastal and Shelf Science* 140, 95-122.
- Ramesh, R., Subramanian, V., (1988). Temporal, spatial and size variation in the sediment transport in the Krishna River basin, India. *Journal of hydrology* 98, 53-65.
- Ranasinghe, P. N., Ortiz, J. D., Moore, A. L., McAdoo, B., Wells, N., Siriwardana, C. H. E. R., Wijesundara, D. T. D. S., 2013. Mid–Late Holocene coastal environmental changes in

- southeastern Sri Lanka: New evidence for sea level variations in southern Bay of Bengal. *Quaternary International* 298, 20-36.
- Sarkar, S.K., Franc̃is̃kovic̃-Bilinski, S., Bhattacharya, A., Saha, M., Bilinski, H., (2004). Levels of elements in the surficial estuarine sediments of the Hugli River, northeast India and their environmental implications. *Environment international* 30, 1089–1098,
- Selvaraj, K., Ram Mohan, V., Szefer, P., (2004). Evaluation of metal contamination in coastal sediments of the Bay of Bengal, India: geochemical and statistical approaches. *Marine pollution bulletin* 49, 174–185.
- Silva, P.N.M.P.D., (2015). Distribution of heavy metals in sediments of Hikkaduwa, Galle, Beruwala fisheries harbors, Sri Lanka (Unpublished thesis). Post Graduate Institute of Science, University of Peradeniya.
- Solali, A., Ghandi, M.S., Kasilingam, K., Sriraman, E., (2013). Heavy Metal Accumulation in the Surface Sediments off Pondicherry, Bay of Bengal, South East Coast of India. *International journal of innovative science, engineering and technology* 2(10), 5741-5753.
- Subramanian V., Jha, P.K., Jergrieken, R., (1988). Heavy metals in the Ganges estuary. *Marine pollution bulletin* 19, 290–293.
- Taylor, S.R., McLennan, S.M., (1985). The continental crust: its composition and evolution. In: *An Examination of the Geochemical Record Preserved in Sedimentary Rocks*. Blackwell scientific publications, Oxford, Edinburgh. 1-30.
- Tomlinson, D. L., Wilson, J. G., Harris, C. R., Jeffrey, D. W., (1980). Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer Meeresuntersuchungen* 33(1), 566.
- Tornero, V., d'Alcalà, M. R., (2014). Contamination by hazardous substances in the Gulf of Naples and nearby coastal areas: a review of sources, environmental levels and potential impacts in the MSFD perspective. *Science of the total environment* 466, 820-840.
- Wang, S. L., Xu, X. R., Sun, Y. X., Liu, J. L., Li, H. B., (2013). Heavy metal pollution in coastal areas of South China: a review. *Marine pollution bulletin* 76(1), 7-15.
- Wang, X.L., Sato, T., Xing, B.S., Tao, S., (2005). Health risks of heavy metals to the general public in Tianjin, China via consumption of vegetables and fish. *Science of the total environment* 350, 28–37.

Young, S. M., Ishiga, H., Roser, B. P., Pitawala, A., (2014). Geochemistry of sediments in three sectors of Trincomalee Bay, Sri Lanka: provenance, modifying factors and present environmental status. *Journal of soils and sediments* 14(1), 204-217.

Zohra, B. S., Habib, A., (2016). Assessment of heavy metal contamination levels and toxicity in sediments and fishes from the Mediterranean Sea (southern coast of Sfax, Tunisia). *Environmental science and pollution research* 23(14), 13954-13963.